

Domino Ant & Cockroach Label

A 3 Pentrich Rd, Masons Mill, Pietermaritzburg, 3201

P +27 (0) 33 342 2111

M info@synvita.co.za

Reg. No. L9782 Act 36 of 1947
Namibian Reg No.: N-AR1794

Read the label before use.
Keep out of reach of children and animals.

GROUP	2B	INSECTICIDES
-------	----	--------------

A scientifically tested ready to use gel bait to control all common species of cockroaches including American, Oriental, and German cockroaches and their nymphs and common species of ants including Black, Sugar and Garden ant.

This insecticidal, non-repellent gel bait is for crack, crevice, and spot treatments in domestic, commercial, and industrial premises. It can be safely used in food handling, storage, and preparation areas.



WARNING !

HAZARD STATEMENTS

Very toxic to aquatic life with long lasting effects.

Precautionary Statements:

Avoid release into the environment.

Collect spillage.

Dispose of contents/container to a licensed hazardous- waste disposal contractor or collection site.

ACTIVE INGREDIENT

Fipronil (phenyl pyrazole) 0,6 g/kg

NET CONTENTS: 35g

Manufactured & Distributed by:

Synvita (Pty) Ltd

Batch no.: See packaging.

Date manufactured: See packaging.

UN NO.: 3082.

In case of poisoning, call the following number. **Emergency number - Griffon Poison Centre 082 446 8946.**

Registration Holder:

Lifeguard Sciences (Pty) Ltd

Reg. No. 2014/079879/07

3 Pentrich Road, Masons Mills, 3201.

+27 33 346 1920.



WARNINGS

- Very toxic to aquatic life with long lasting effects.
- Handle with care.
- Keep out of reach of children, uninformed persons, and animals.
- Store in the closed, original container in a cool place and out of direct sunlight.
- Do not break open the DOMINO Ant & Cockroach Gel syringe.

NOTICE TO THE USER: DOMINO Ant & Cockroach Gel is to be used only in accordance with the instruction on the label. It is an offense under the Act to use this product in a manner contrary with the directions on the label.

PRECAUTIONS

- Avoid contact with eyes and skin.
- Do not eat drink or smoke when applying bait.
- Wash hands with soap and water after use.
- Avoid contamination of food, feed, water, eating utensils and food preparation surfaces.
- Never use the empty container for any other purpose.
- Do not place the DOMINO Ant & Cockroach Gel on any hot surface, in the rain or in direct sunlight.
- Triple rinse empty pesticide containers or packages by triple rinsing before disposal. Quarter-fill the empty packaging with water, close and shake for 30 seconds. Empty the content by placing it upside down over disposal bucket.
- Avoid release into the environment.
- Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site.

RELEVANT SUBSTANCES

Fipronil tech, <0.1%, Cas No.: 120068-37-3; Sapogenat, <1%, Cas No.: 9046-09-7

FIRST AID

Provide medical personnel with the container label, product name or safety data sheet (SDS) when seeking medical attention.

IF IN EYES: Rinse cautiously with water for several minutes (15minutes). Remove contact lenses, if present and easy to do. Continue rinsing.

IF INHALED: Remove person to fresh air and keep comfortable for breathing.

IF ON SKIN: Wash thoroughly with soap and water. Do not rub skin hard.

IF SWALLOWED: Keep patient calm and reassured, rinse mouth with water.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

AREA OF APPLICATION:

DOMINO Ant & Cockroach Gel may be used both internally and externally. Direct sunlight and water may reduce the residual effect.

- Domestic premises including kitchens and bathrooms.
- Commercial kitchens, canteens, restaurants, food preparation, processing, and storage premises.
- Public Buildings including retail outlets, hospitals, hotels, schools, churches.
- Commercial and industrial premises including factories, shops, and warehousing.
- Aircraft, motor vehicles, railway carriages, ships, and boats.

PEST	LOCATIONS	RATE OF APPLICATION	COMMENTS
Common cockroaches including German, Oriental and American cockroach and their nymphs and common species of ants including Black ant, Common ant, Sugar ant and Garden ant.	Domestic, Commercial, and Industrial premises including food handling, storage, and preparation areas.	Apply 1-3 spots per m ² . Recommended spot size is 0,03g – 0,06g. Depending on the severity of the infestation and species present it may be necessary to increase the number of spots to three and increase the spot size.	Placements of spots should be in areas where cockroaches and ants are known to harbour, such as crack, crevice, etc. (See Area of Application above.)

GENERAL APPLICATION INSTRUCTIONS

- Remove cap and place the tip on the surface to be treated, depress plunger slightly.
- High cockroach and ant challenge may require ribbon strips to be applied.
- Recap after use and place in its outer pack.
- Re-apply according to level of infestation.
- A visual inspection of gel placement is recommended two weeks to one month after initial treatment.