

Pulse Wax Blocks - Label

A 3 Pentrich Rd, Masons Mill, Pietermaritzburg, 3201

P +27 (0) 33 342 2111

M info@synvita.co.za

Reg. No. L6189 Act 36 of 1947

Namibian Reg No.: N-AR1213

READ THE LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Rapid Single-Feed.

A highly active anti-coagulant bait for the control of the Norway rat, Roof rat, the House mouse and Gerbil.
Effective against warfarin resistant rats and mice.



HARMFUL

HAZARD STATEMENTS:

Causes damage to organs through prolonged or repeated exposure (blood).

PRECAUTIONARY STATEMENTS:

Do not breathe dust.

Wear protective gloves.

Get medical help if you feel unwell.

ACTIVE INGREDIENT:

Brodifacoum (anti-coagulant): 0.05g ai/kg (0.005%)

NET VOLUME: 80g / 500g/ 3kg/ 10kg

MANUFACTURED & DISTRIBUTED BY: Synvita (Pty) Ltd

REGISTRATION HOLDER: Lifeguard Sciences (Pty) Ltd

Reg. No. 2014/079879/07

3 Pentrich Road, Masons Mill, Pietermaritzburg, 3201.

Tel.: 033 346 1920

Emergency number of Griffon Poison Centre: 082 446 8946

Emergency number of National Poison Control Centre: 0861 555 777

Batch No: See Container

Date manufactured: See Container

UN No.: Not regulated

THIS REMEDY CONTAINS AN ANTI-COAGULANT THAT MAY CAUSE INTERNAL BLEEDING IF CONSUMED. THIS REMEDY MAY ONLY BE USED IN TAMPER-RESISTANT BAIT STATIONS AND ONLY AS DIRECTED ON THIS LABEL. OBTAIN, READ AND FOLLOW ALL SAFETY INSTRUCTIONS BEFORE USE.

READ ATTACHED PACKAGE LABEL BEFORE USE

WARNINGS

- Causes damage to organs through prolonged or repeated exposure (blood).
- Ensure the re-closable lid is tightly closed at all times.
- Immediately after use, store the container with unused bait in its original container, locked up.
- This remedy may ONLY be used in tamper-resistant bait boxes.
- In case of poisoning or suspected poisoning, especially involving children, get immediate medical attention and make this label available.
- Always make sure that baits are adequately protected from children and non-target animals.

PRECAUTIONS

- Obtain, read and follow all safety instructions before use.
- Wear protective gloves.
- Do not breathe dust.
- Get medical help if you feel unwell.
- Store locked up.
- Dispose of contents/container in a responsible manner to avoid accidental exposure of uninformed persons and animals.
- Triple rinse empty pesticide containers or packages by triple rinsing before disposal. Quarter-fill the empty packaging with water, close and shake for 30 seconds. Empty the content by placing it upside down over disposal bucket.

RELEVANT SUBSTANCES: Not Regulated

FIRST AID TREATMENT

If in Eyes: Immediately wash eyes with copious amounts of clean water for at least 15 minutes.

Check for and remove any contact lenses. Continue rinsing. If eye irritation persists, get medical help.

If Inhaled: Remove patient from source of poisoning to a well-ventilated area and loosen constricting clothes. If not breathing, give artificial breathing. If breathing is difficult, give oxygen. Get medical attention immediately.

If on Skin: Immediately remove contaminated clothing, rinse affected skin area with plenty of cool water for at least 15 minutes. Wash clothing before reuse. Thoroughly clean shoes before reuse. Get medical attention immediately.

If Swallowed: Keep patient calm and reassured, rinse mouth with water and ensure airway is clear. Do not induce vomiting unless directed to do so by medical personnel. Never give anything by mouth to an unconscious person. Loosen tight clothing such as a collar, tie, belt or waistband. Get medical attention immediately.

MODE OF ACTION

Brodifacoum is a second-generation anticoagulant that disrupts the vitamin K cycle by inhibiting Vitamin K epoxide reductase, reducing the ability of rodents' blood to clot over time.

SYMPTOMS OF POISONING

Bleeding from the gums and nose, bleeding under the skin, pallor and haemorrhage into and around elbow and knee joints. Blood in urine and faeces.

NOTE TO PHYSICIAN

Intravenous Vitamin K1, 50-150mg on first day, in divided doses, depending on prothrombin time. Keep under medical supervision until prothrombin times return to normal. In severe cases small transfusions of fresh blood.

NOTICE TO USER: Pulse Wax Blocks is to be used only in accordance with the instruction on the label. It is an offense under the Act to use this product in a manner contrary with the directions on the label.



DIRECTIONS FOR USE

USE ONLY AS DIRECTED:

For use in the home, on the farm and industrial premises. **Pulse Wax Blocks** can kill rats and mice after a single feed. Death of rodents will take 4-12 days. Effective against warfarin resistant rats and mice.

NORWAY AND ROOF RATS:

Place **Pulse Wax Blocks** in a covered bait station to prevent access by children and domestic animals.

Place 50-70g of **Pulse Wax Blocks** per bait stations along the walls and active rat runs, in rat burrows and in sheltered areas where rats are known to live or feed. Bait points should be inspected and replenished every 7 days until feeding ceases.

MICE:

Place 10-20g **Pulse Wax Blocks** per bait station every 2-4m along the wall or runways, in nests and in sheltered areas where mice are known to live and feed. The single feeding action of **Pulse Wax Blocks** is particularly effective against mice with their small appetites and sporadic feeding behaviour.

GERBIL:

Place 50-70g **Pulse Wax Blocks** in protected bait stations 10m apart or 10-20g placed into active holes in the colony.